

Kettlethorpe


HIGH SCHOOL

MATHS

Year 10 | Theta

Name:

Set:



Unit	Topic	Complete
1	Percentages	
2	Multiplicative Reasoning	
3	Averages	
4	Representing Data	
5	Scatter Graphs	
6	Linear Graphs	
7	Quadratic, Cubic and Other Graphs	
8	Real Life Graphs	
9	Inequalities	
10	Perimeter and Area	
11	3D Shapes	
12	Constructions and Loci	
13	Similarity and Congruence	

Theta Unit 1: Percentages

Prior Knowledge

Convert between fractions decimals and percentages.

Express a given number as a percentage of another number, including where the percentage is greater than 100%.

E.G.

Express $\frac{57}{60}$ as a percentage $\frac{57}{60} = \frac{19}{20} = \frac{95}{100} = 95\%$.

Calculate percentage change with or without a multiplier.

E.G.

Increase 70 by 30%

Without multiplier, find $30\% = 21$ add this to original $70 + 21 = 91$.

With a multiplier $100\% + 30\% = 130\% = 1.3$
 $70 \times 1.3 = 91$.

Find an original quantity after a percentage change.

E.G.

The cost of a ticket has been increased by 12.5% to £225, find the original amount.
 $£225 \div 1.125 = £200$.

Compound Percentages

Example

James invests £6000 for 5 years, at 3% a year.
At the end of 5 years, calculate the value of the investment after 5 years.

Answer:

$$100\% + 3\% = 103\% = 1.03$$

$$6000 \times 1.03^5$$

$$£6955.64$$

Example 2

Katy invests £2000 for 3 years.
She receives interest per year of 2.5% in the first year then $x\%$ for the second and third year.
There is a total of £2124.46 at the end of the 3 years. Work out the value of x .

Answer:

Let y be the multiplier for the second and third year.

$$2000 \times 1.025 \times y^2 = 2124.46$$

$$y = \sqrt{\frac{2124.46}{2000 \times 1.025}}$$

$$y = 1.017999$$

Therefore the increase, x , is 1.8%.

Literacy

Explain how you would find the multiplier to work out a 2.5% decrease.

Reasoning

Chloe is given a 10% pay rise. The next year Chloe is given another 10% pay rise. Her manager says that Chloe's pay has increased by 20% overall. Explain why Chloe's manager is wrong.

Fluency

Percentages of amounts non calculator

- 1) Find 35% of 160 2) Increase £140 by 15%

- 4) The cost of a shirt was reduced by 20% in a sale. The shirt costs £56 in the sale. What was its original price?

Percentages of amounts, calculator allowed, use multipliers to

- 1) Increase £270 by 6.5% 2) Decrease 180 kg by 3.5%

- 4) The cost of a laptop was reduced by 35% in a sale. The laptop costs £325 in the sale. What was its original price?

Problem Solving

James invests £5000 for 4 years, at 3% a year. Work out the value of the investment after 4 years.

Theta Unit 2: Multiplicative Reasoning

Prior Knowledge

Use simple unitary proportion.

Understand scaling recipes.

Rearrange formulae.

Solve equations.

Work out best buys.

Fractions and Ratio

The denominator of the fraction is the total number of parts added together.

$$4 : 3 = \frac{4}{7} : \frac{3}{7}$$


4 + 3 = 7

Best Buys

E.G.



Pack of 4
toilet rolls
£1.96



Pack of 9
toilet rolls
£4.23

Which pack offers the best value for money?

We need to work out the cost for 1 toilet roll from each.

Option 1 (9 rolls)
1 toilet roll costs
 $£4.23 \div 9 = 47p$.

Option 2 (4 rolls)
1 toilet roll costs
 $£1.96 \div 4 = 49p$.

Conclusion:
Option 1 is better value.

You must always show your working on these questions.

Simple Compound measures

You must know and be able to use the formulas and be able to rearrange them when needed.

$$\text{speed} = \frac{\text{distance}}{\text{time}} \quad \text{density} = \frac{\text{mass}}{\text{volume}}$$

$$\text{pressure} = \frac{\text{force}}{\text{area}}$$

You will also need to be able to convert between their measures, i.e. convert between mph and miles per minute.

Harder Compound measures

Use a table to help you organise your working out. This is helpful when there are multiple parts to journeys.

Example:

Sienna travels from Birmingham to Leeds at an average speed of 60mph.

She then travels from Leeds to Darlington at an average speed of 40mph.

The distance from Birmingham to Leeds is 150miles.

The distance from Leeds to Darlington is 70 miles.

Calculate Sienna's average speed.

	B TO L	L TO D	TOTAL B TO D
SPEED	60mph	40mph	51.76mph
DISTANCE	150miles	70miles	220miles
TIME	2.5h	1.75h	4.25h

Literacy

Write the definition of density.

Write the definition of mass.

Reasoning

If wood has density less than 1g/cm^3 it will float.

Which of these will be best for building a toy boat?

Plank A

Volume = 750cm^3
Mass = 900g

Plank B

Volume = 0.0152m^3
Mass = 7.6kg


Plank C

Volume = 1000cm^3
Mass = 1.02kg

Fluency

The distance from Caxby to Drone is 45 miles.

The distance from Drone to Elton is 20 miles.



Colin drives from Caxby to Drone. Then he drives from Drone to Elton.


Colin drives from Caxby to Drone at an average speed of 30 mph.

He drives from Drone to Elton at an average speed of 40 mph.

Work out Colin's average speed for the whole journey from Caxby to Elton.

Problem Solving

The diagram below shows a solid block of ice.



81cm

A block of ice weighs $\frac{1}{2}$ tonne.

The block is a cube with side length 81cm.

Find the density of the ice.

Give your answer in kilograms per cubic metre.

Theta Unit 3: Averages

Prior Knowledge

Understand the difference between quantitative and qualitative data.

Understand the difference between continuous and discrete data.

Put data into a frequency table.

Put data into a grouped frequency table.

Find the averages, mode, mean and median for a data set.

Find the range for a data set.

Compare data using an average and range.

Median and Mode from frequency table

Here is a table showing the number of goals scored in 10 football matches.

Number of goals	Frequency
0	2
1	2
2	5
3	1

Mode = 2 (the class with highest frequency)

The **median** is the class containing the 5.5th data point.

Number of goals	Frequency	Cumulative
0	2	2
1	2	2+2 = 4
2	5	4+5 = 9
3	1	9+1 = 10

The 5.5th data point is in the category for 2, therefore the median is 2.

Mean from Frequency Table

To find the mean, you need to find the total number of goals scored.

Number of goals, g	Frequency, f	g x f
0	2	0
1	2	2
2	5	10
3	1	3

Total goals $0 + 2 + 10 + 3 = 15$.

Mean = $15/10 = 1.5$ goals per game.

Remember

When the data is grouped like below, we estimate the mean using the midpoint for the classes.

Length (l, cm)	Midpoint (M)	Frequency, f	M x f
$10 < l \leq 20$	15	10	150
$20 < l \leq 40$	30	30	900
$40 < l \leq 50$	45	20	900

Estimated total length = $150 + 900 + 900 = 1950$.

Estimated Mean = $1950 \div 60 = 32.5$ goals per game.

Literacy

Explain the meaning of quantitative and qualitative data.

Reasoning

Decide if the statements are true or false. Give a reason for each of your answers.

Score	Frequency
11	3
12	3
13	2
14	3
15	1

A The data set contains exactly 10 values.

B The median is 13 because it's in the middle of: 11, 12, 13, 14, 15.

C The mean can't be 12.7 because only whole numbers could be scored.

D The mean score is 2.4.

E The data is discrete.

F The modal value is 3.

Fluency

Calculate the mean, median and mode from each table.

(a)

Age	Frequency
5	2
6	2
7	5
8	1

(b)

Number of phones	Frequency
0	1
1	3
2	2
3	0
4	4
5	0

(c)

Mass	Frequency
$20 < m \leq 25$	12
$25 < m \leq 30$	24
$30 < m \leq 35$	17
$35 < m \leq 40$	15
$40 < m \leq 45$	4

(d)

Height	Frequency
$120 < h \leq 130$	51
$130 < h \leq 140$	120
$140 < h \leq 150$	66
$150 < h \leq 160$	59
$160 < h \leq 170$	4

Problem Solving

Test Scores		
	BOYS	GIRLS
Mean	32 marks	40 marks
Range	18	15

Compare the distributions of boys and girls test scores.

Theta Unit 4: Representing Data

Prior Knowledge

Understand the difference between quantitative and qualitative data.

Understand the difference between continuous and discrete data.

Compare data using an average and range.

Draw and interpret bar charts.

Draw and interpret composite and dual bar charts.


Draw and interpret pie charts.

Frequency Polygons

- Plot midpoints.
- Join dot-to-dot.

E.G.

Number of Sweets	Mid Value	Frequency
12 - 16	14	8
17 - 21	19	11
22 - 26	24	19
27 - 31	29	16
32 - 36	34	5




Constructing a Histogram

Histograms are used to represent continuous data. We plot bars that are the width of the classes, but for the height, we use frequency density, which is given by the formula:

$$\text{Frequency density} = \frac{\text{frequency}}{\text{class width}}$$

Example

HEIGHT (CM)	FREQUENCY	CLASS WIDTH	FREQUENCY DENSITY
65 < h ≤ 75	2	10	2/10 = 0.2
75 < h ≤ 80	7	5	7/5 = 1.4
80 < h ≤ 90	21	10	21/10 = 2.1
90 < h ≤ 105	15	15	15/15 = 1
105 < h ≤ 110	12	5	12/5 = 2.4



Angles for a Pie Chart:

Step 1: Calculate total of frequencies.

Step 2: $360 \div \text{Total} = a$

Step 3: Multiply a by each frequency.

Mobile Phone	Frequency	
Apple	12	$\times 10 = 120^\circ$
Samsung	9	$\times 10 = 90^\circ$
HTC	8	$\times 10 = 80^\circ$
Sony	6	$\times 10 = 60^\circ$
Nokia	1	$\times 10 = 10^\circ$

Literacy

Explain the meaning of discrete and continuous data.


Reasoning

Dahlia says “if a bar on a histogram is twice as tall as another bar, then it represents twice as many people”
Is she correct? Give a reason for your answer.

Fluency


The table gives information about the money, £ A , some people spent on an internet site one day. On the grid, draw a frequency polygon for this information.

Money spent (£ A)	Frequency
$0 < A \leq 20$	10
$20 < A \leq 40$	15
$40 < A \leq 60$	25
$60 < A \leq 80$	40
$80 < A \leq 100$	6



Problem Solving

The pie chart shows types of properties in a street.
There are 80 Bungalows.



Work out the number of semi detached properties.

Theta Unit 5: Scatter Graphs

Prior Knowledge

Plot coordinates.

Read diagrams.

Plot a scatter graph.




Interpreting Correlation

A frequency polygon is used to plot grouped data, it is plotted as mid-point against frequency, the points are then joined using a ruler.

Positive correlation means as one variable increases so does the other variable.

Negative correlation means as one variable increases the other variable decreases.

No correlation means there is no relationship between the two variables.



Interpolation and Extrapolation


Scatter diagrams can be used to make estimates, first we need to draw a line of best fit.

Example:

For the data in the example, estimate how many umbrellas would be sold when there is 3mm of rain.

Therefore reading from our line of best fit, we'd estimate 19 umbrellas to be sold. This is **interpolation** as it is within the data range we have.

When we are asked to do this outside of the data range it is called **Extrapolation** and is less accurate as we have no data in this range.



Literacy

Explain what is meant by extrapolation and interpolation.

Fluency Reasoning

The scatter graph shows information about 10 adult snakes of the same type.

It shows the length and weight of each snake.

An adult snake of this type has a weight of 740 g.

(a) Use the scatter graph to estimate the length of this snake.

Another snake measured 68cm and had a weight of 848g


(b) Show this information on the scatter graph.

(c) This snake is an outlier, give a possible reason for this.

Steven wants to estimate the weight of an adult snake of length 110 cm.

He says he will draw a line of best fit and read off the weight at 110 cm.

(b) Explain what is wrong with his method.



Theta Unit 6: Linear Graphs

Prior Knowledge

Plot Linear graphs.

Identify the gradient, m , of a linear graph.

Find the equation of a straight-line graph in the form $y=mx + c$.

Identify parallel lines, they have the same gradient.

Identify perpendicular lines, the gradients are the negative reciprocal of one another.

Equation of a Straight Line

The equation of a line is in the form:

$$y = mx + c$$


M is the gradient. C is the y intercept

E.G.

$$m = 3 \div 1 = 3$$

$$c = -6 \text{ (crosses } y\text{-axis)}$$

$$\text{So } y = 3x - 6$$




Equation of a Line Given Two Points

To find the equation of a line between two points, first it's good to draw a picture, find the gradient between the two points, then use this in the equation for m and use one point to find the intercept.

Example

Find the equation of the line that passes through the points $(3,5)$ and $(7, 19)$



$$\text{Gradient} = \frac{17-5}{7-3} = \frac{12}{4} = 3$$

+ c

$$5 = 3 \times 3 + c$$

$$5 = 9 + c$$

$$\text{so } y = 3x$$

$$\text{Answer } y = 3x - 4$$

Literacy

Give the definitions of the words parallel and perpendicular.

Reasoning

A line has the equation $y = 2x - 3$.



Student A says: the perpendicular gradient is -2

Student B Says: the perpendicular gradient is $\frac{1}{2}$

Comment on their answers

Fluency

Find the equations of these lines.



Find the equation of the line that passes through the points $(4, 7)$ and $(6, 15)$

Write down the equation of each of the following line parallel to $y = 3x + 5$ and passing through $(0, 2)$

Write down the equation of each of the following line parallel to $y = 2x + 4$ and passing through $(0, 3)$

Problem Solving

Do the points $(1, 4)$, $(4, 10)$ and $(9, 20)$ lie in a straight line?

Theta Unit 7: Quadratic, Cubic and Other Graphs

Prior Knowledge


Draw linear graphs.

Solve linear equations.

Substitute into quadratic expressions.

Complete a table of values for and plot a quadratic graph.
E.G. Plot the graph of $y = x^2 - x - 6$


x	y
-3	6
-2	0
-1	-4
0	-6
1	-6
2	-4
3	0
4	6



Key Graphs


There are key graphs that you need to be able to recognise and know the general form.

Exponential




$$y = a^x$$

Reciprocal




$$y = \frac{a}{x}$$

Cubic



$$y = x^3$$

Quadratic



$$y = x^2$$


Solutions from Quadratic Graphs

This is where the graph crosses the x axis. They can be called solutions or roots.

Example

Given the graph $y = x^2 + x - 3$.

Roots: $x = -2.3$ or $x = 1.3$.









Literacy

Explain what roots of an equation means.

Fluency

Here are six graphs. Give the name of each type of graph.



Write down the letter of the graph that could have the equation

1) $y = 2^x$

2) $y = x^3 - 3x$

3) $y = 3x$

4) $y = x^3 - 3x$


5) $y = x^2 - 5$

Reasoning Problem Solving

Complete the table of values for $y = 2x^2 - 1$

x	-2	-1	0	1	2
y	7			1	

On the grid below, draw the graph of $y = 2x^2 - 1$ for values of x from $x = -2$ to $x = 2$



Use your graph to write down estimates of the solutions of the equation $2x^2 - 1 = 0$

Theta Unit 8: Real Life Graphs

Prior Knowledge

Plot coordinates in all 4 quadrants.

Find the mid-point of a line segment.

Use Pythagoras on right-angle triangles.

Use a conversion graph.


Draw and interpret distance-time graphs.

Distance Between Two Points

The distance between two points, can be seen as a right angle triangle, so we can use Pythagoras to find the distance between two points.

E.G. Find the distance between the points (3,2) and (8,9).

$$\sqrt{7^2 + 5^2} = \sqrt{74} = 8.60232 \dots$$



Real Life Graphs

Example

The graph shows the cost of hiring a chainsaw from saws r us. The company charge a fixed charge plus a daily charge.

Calculate the fixed and daily charge.


Answer:

Fixed Charge = Starting price for 0 days the intercept, this is £10.

Daily charge = This can be calculated by working out the difference from one day to the next. Day 0 = £10, Day 2 = £12.50.

$$12.50 - 10 = 2.50 \quad 2.50 \div 2 = £1.25$$

Daily charge = £1.25.



Literacy

Give the definitions of speed and velocity.

Reasoning

A, B and C are three points.

Point A has coordinates $(-5, 4)$.

Point B has coordinates $(0, 10)$.

Point B is the midpoint of A and C.

Sam says Point C has coordinates $(-2.5, 7)$

Explain Sam's mistake and give the correct coordinates for Point C.

Fluency



Find the mid-point of each pair of coordinate.

1) $(2, 3)$ and $(0, 11)$
 $-2)$

2) $(5, 10)$ and $(-1, 12)$


3) $(-1, 5)$ and $(-2, -2)$

Calculate the distance between the two points.



Problem Solving

The graph shows the cost of hiring a rotator from tools r us.



The company charge a fixed charge plus a daily charge.

Calculate the fixed and daily charge

Theta Unit 9: Inequalities

Prior Knowledge

Know what integer means (whole number).

Use inequality notation:

< means less than
 \leq means less than or equal to
 > Means more than
 \geq means more than or equal to


Solve linear equations.

Substitute into expressions.


Factorise expressions.

Representing Inequalities on a Number Line

We can represent inequalities on a number line, hollow means not included, filled in means included.




$x < 2$, x is less than 2



$x > 2$ x is greater than 2



$x \leq 2$ x is less than or equal to 2



$x \geq 2$ x is greater than or equal to 2

Solving Inequalities

Sometimes we solve linear inequalities, which are solved the same as equations using balancing method.

E.G.

Solve $5x - 24 > 11$

$$\begin{array}{rcl} 5x - 24 & > & 11 \\ +24 & +24 & \\ \hline 5x & > & 35 \\ \div 5 & \div 5 & \\ \hline x & > & 7 \end{array}$$

Solving inequalities gives a range of answers, rather than an individual solution.

Solving Two Linear Inequalities

Sometimes we have a group of two inequalities, you still use the balancing method, doing the same inverse operation to all 3 parts of the inequality.

E.G.

Give the integers $3 < 2x - 5 \leq 11$

$$\begin{array}{rcl} 2 < 2x - 5 \leq 11 \\ +5 & +5 & +5 \\ \hline 7 < 2x \leq 16 \\ \div 2 & \div 2 & \div 2 \\ \hline 3.5 < x \leq 8 \end{array}$$

The integers (whole numbers) which satisfy this inequality are:

4, 5, 6, 7, 8.

Literacy

Write the definition of Inequality

Use the word inequality within a sentence

Reasoning

Solve $7 - 4x > 15$

Ama gives the following solution

$$11 - 8x > 15 \quad (-7)$$

$$8x > 4 \quad (\div 8)$$

$$x > 2$$

Comment on Ama's solution.

Fluency

List the integer solutions for the following:

1) $-2 < x \leq 5$ 2) $4 \geq x > -4$ 3) $-3 \leq 2x \leq 10$

Solve the following:

1) $2x + 7 \leq 15$ 2) $4x + 13 \geq 6$ 3) $5x - 2 \geq 2x + 13$

4) $2(x + 7) \leq 5(2x + 3)$ 5) $7 - 3x < 19$ 6) $12 - 2x > 8 - 5x$

Problem Solving

1) Given that a and b are integers such that

$$10 < 2a < 22$$

$$-3 < b < 6$$

and $a + b = 9$

Find all the possible values of a .

2) Bianca, Bob and Valentina have completed some Hegarty tasks. Bianca has completed 40 more than Bob. Valentina has completed 3 times as many as Bob.
Together Bob and Bianca have completed twice as many as Valentina.
Calculate the least number of tasks Bob could have completed.

Theta Unit 10: Perimeter and Area

Prior Knowledge

Perimeter is the distance around the outside of a 2D shape.

Area is the amount of space inside a 2D shape.

Identify the names for parts of a circle.

Calculate the area of basic and compound shapes.

Convert between metric measures for area.

Know the following key formulae:

Area triangle = $\frac{1}{2} \times \text{base} \times \text{perpendicular height}$

Area parallelogram = $\text{base} \times \text{perpendicular height}$

Area trapezium
= $\frac{1}{2}(a + b) \times \text{height}$

Area of a circle = πr^2

Circumference of circle = πd

Area of Sector

The area of a sector is calculated by the formula:


$$\frac{\theta}{360} \times \pi r^2,$$

where θ = the angle inside the sector.

Example

Find the area of the sector enclosed by two radii of 4cm and 60° .

$$\text{Area} = \frac{60}{360} \times \pi \times 4^2 = 8.4\text{cm}^2$$



Arc Length


The arc length is calculated by the formula $\frac{\theta}{360} \times \pi 2r$ or $\frac{\theta}{360} \times \pi d$,

where θ = the angle inside the sector.

Example

Find the minor arc length, enclosed by radii of 3.5cm and 144° .

$$\text{Arc length} = \frac{144}{360} \times \pi \times 2 \times 3.5 = 8.8\text{cm}$$



Literacy

Write the definition of a Trapezium.

Use the word trapezium within a sentence.


Reasoning

Sameera says that you cannot draw a square that has a perimeter of x cm and an area of $x \text{ cm}^2$ (where x is the same value in each case). Is Sameera correct? Give a reason for your answer. .


Fluency

Find the area of the following shapes:.

1)



2)




Find the area of a sector with radius 8 cm and angle between the two radii of 82° .

Find the arc length of a sector with radius 4.6 cm and an angle between the two radii of 145° .

Problem Solving

1) A circular pie is cut in to 8 slices of equal shape. The area of the top of one of these slices is 48 cm^2 . Find the diameter of the pie.

2) The area of this right-angled trapezium is 26 m. Find the perimeter of the trapezium.



Theta Unit 11: 3D Shapes

Prior Knowledge

Volume is the space inside a 3D shape.

Calculate the volume of a prism, including cylinders.

Draw plans and elevations for shapes.

Convert between metric measurements for volume.

Calculate surface area of 3D shapes.

Spheres

If required to calculate the volume or surface area of a sphere then the following formulae will be given.

$$\text{Volume of sphere} = \frac{4}{3} \times \pi \times r^3$$

$$\text{Surface area of sphere} = 4 \times \pi \times r^2$$

Volume of Pyramid


The formula volume of a pyramid will not be given to you. The volume is calculated from the formula.

$$\frac{1}{3} \times \text{area of base} \times \text{height}$$

Remember: a cone is a circular based pyramid

Curved Surface Area of Cones


For the curved surface area of a cone the formula is $\pi \times r \times l$



Remember: for total surface area add the area of the base on.

Frustums

A frustum is a truncated pyramid, to calculate the volume, take the volume of the smaller cone away from the volume of the larger cone.



Literacy

Write the definition of prism.

Use the word prism within a sentence.

Reasoning


Jamal says "if a cone and a sphere have the same volume and also the same radius, then the height of the cone is bigger than the radius". Is he correct? How much more is the height compared to the radius?

Fluency

Find the volume and surface area of the following shapes:


- 1) A sphere with **diameter** of 9m and height m.
- 2) A cone with radius of 3.8cm and height 9cm.

- 3) Calculate the total volume of this 3D shape.



Problem Solving

The conical shaped container is partially filled with water, so that the water has depth of 9cm. What is the volume of the water in the container?



Theta Unit 12: Constructions and Loci


Prior Knowledge

Using a compass. Using a protractor. Draw and use Loci. Draw plans and elevations of 3D shapes.
Construct a perpendicular bisector. Construct an angle bisector. Estimate lengths from scale diagrams.
Measure and use bearings. Know that the perpendicular from a point to a line is the shortest distance to the line.
Draw triangles accurately using a protractor and compass.

Using Loci to Find Regions on Scale Diagrams

Jane wants to plant a tree in the garden. It needs to be at least 5m from C. Nearer to AB than AD and less than 3m from DC.

On the diagram, shade the region where Jane should plant the tree.



Arc radius around C.

Bisector of AB and AD to find points nearer AB than AD.

Line as a loci from DC to show close enough to DC.


Region shaded red.

Bearings and Loci

Here is a map. S is the position of a ship. C is a point on the coast.

Ships must not sail within 500m of C.
The ship is on a bearing of 037° .

Will the ship sail closer than 500m of C?



Circle around C to represent the area close enough to the ship.

Then the bearing is shown as well.

Yes, as the ship's course does intersect the region.

Literacy

Write the definition of loci.

Use the word loci within a sentence.

Reasoning


Nathan was asked to show all the points equidistant from A and B. Comment on his response.



Fluency

Find the following bearings

- 1) The bearing of A from B is 138° . What is the bearing of B from A?
- 2) The bearing of C from D is 284° . What is the bearing of D from C?
- 3) The bearing of E from F is 082° . What is the bearing of F from E?



Label the region R,
that is:

Closer to AB than CD

Closer to CD than AD

More than 3cm from C

Problem Solving

Ship A sails on a bearing of 060° at 25km an hour.
Ship B sails on a bearing of 285° at 40km an hour.
If both boats set off at the same time, how far apart are they after 4 hours? Use a scale of 1cm = 30km



Theta Unit 13: Similarity and Congruency

Prior Knowledge

Understand ratios.

Know angle rules.

Construct a geometric argument with angles.

Understand properties of triangles.

Know that enlarging a shapes sides by scale factor 2, doesn't increase the area by scale factor 2.


Apply similarity to 2D shapes to find missing sides.

Area Scale factor

The area scale factor for 2 shapes is the square of the linear scale factor.

E.G.

Quadrilaterals P and Q are similar, the area of P is 10cm^2 . Calculate the area of Q.



$$LSF = \frac{12}{3} = 4$$

$$ASF = 4^2 = 16$$


$$\text{New area} = 16 \times 10 = 160\text{cm}^2$$

Volume scale factor

The volume scale factor for 2 shapes is the cube of the linear scale factor.

E.G.

Quadrilaterals R and S are similar, the volume of R is 40cm^3 . Calculate the volume of S.



$$LSF = \frac{35}{14} = 2.5$$

$$VSF = 2.5^3 = 15.265$$


$$\text{New volume} = 40 \times 15.265 = 625\text{cm}^3$$

Congruency

In order to prove 2 triangles are congruent they must share 3 pieces of information.

A proof must contain 3 bullet points, each stating the link and why they are the same.

Reasons can be given in the question or by shared side.



Literacy

Write the definition of similar.

Use the word similar within a sentence.

Reasoning


A and B are similar cubes. The length of each edge on cube A is

y cm and the length of each edge on cube B is $2y$ cm.

Ken says "Everything about Cube B is twice as big as Cube A"

Comment on Ken's statement.

Fluency




Work out the length of FG given these are similar rectangles.

A and B are similar solids.

The surface area of shape B is 400cm^2 .

Find the surface area of A.



Problem Solving

Chocopuffs are sold in small and large boxes.

The boxes are similar cuboids.

Volume of A : Volume of B = 8 : 125

The front of the larger box has an area of 500cm^2 .

What is the area of the front of the smaller box?